§ 866.3700

§866.3700 Staphylococcus aureus serological reagents.

Identification. Staphylococcus aureus serological reagents are devices that consist of antigens and antisera used in serological tests to identify enterotoxin (toxin affecting the intestine) producing staphylococci from cultured isolates. The identification aids in the diagnosis of disease caused by this bacterium belonging to the genus Staphylococcus and provides epidemiological information on these diseases. Certain strains of Staphylococcus aureus produce an enterotoxin while growing in meat, dairy, or bakery products. After ingestion, this enterotoxin is absorbed in the gut and causes destrucof the intestinal tion (gastroenteritis).

(b) Classification. Class I (general controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to the limitations in §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 54 FR 25047, June 12, 1989; 66 FR 38792, July 25, 2001]

§866.3720 Streptococcus spp. exoenzyme reagents.

(a) Identification. Streptococcus spp. exoenzyme reagents are devices used to identify antibodies to Streptococcus spp. exoenzyme in serum. The identification aids in the diagnosis of disease caused by bacteria belonging to the genus Streptococcus and provides epidemiological information on these diseases. Pathogenic streptococci are associated with infections, such as sore throat, impetigo (an infection characterized by small pustules on the skin), urinary tract infections, rheumatic fever, and kidney disease.

(b) Classification. Class I (general controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to the limitations in §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 61 FR 1119, Jan. 16, 1996; 66 FR 38792, July 25, 2001]

§866.3740 Streptococcus spp. serological reagents.

(a) *Identification*. *Streptococcus* spp. serological reagents are devices that

consist of antigens and antisera (excluding streptococcal exoenzyme reagents made from enzymes secreted by streptococci) used in serological tests to identify Streptococcus spp. from cultured isolates derived from clinical specimens. The identification aids in the diagnosis of diseases caused by bacteria belonging to the genus Streptococcus and provides epidemiological information on these diseases. Pathogenic streptococci are associated with infections, such as sore throat, impetigo (an infection characterized by small pustules on the skin), urinary tract infections, rheumatic fever, and kidney disease.

(b) Classification. Class I (general controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 65 FR 2312, Jan. 14, 2000]

§866.3780 Toxoplasma gondii serological reagents.

(a) Identification. Toxoplasma gondii serological reagents are devices that consist of antigens and antisera used in serological tests to identify antibodies to Toxoplasma gondii in serum. Additionally, some of these reagents consist of antisera conjugated with a fluorescent dye (immunofluorescent reagents) used to identify Toxoplasma gondii from clinical specimens. The identification aids in the diagnosis of toxoplasmosis caused by the parasitic protozoan Toxoplasma gondii and provides epidemiological information on this disease. Congenital toxoplasmosis is characterized by lesions of the central nervous system, which if undetected and untreated may lead to brain defects, blindness, and death of an unborn fetus. The disease is characterized in children by inflammation of the brain and spinal cord.

(b) Classification. Class II (performance standards).

\$ 866.3820 Treponema pallidum non-treponemal test reagents.

(a) Identification. Treponema pallidum nontreponemal test reagents are devices that consist of antigens derived from nontreponemal sources (sources not directly associated with